

Trends in dental caries in 12- and 13-year-old schoolchildren from Florianópolis between 1971 and 2009

Helena Mendes Constante¹, João Luiz Bastos², Marco Aurélio Peres²

¹ Undergraduate dental student, Federal University of Santa Catarina, Brazil

² DDS, MSc, PhD, Professor, Department of Public Health, Federal University of Santa Catarina, Brazil

Abstract

Aim: To estimate the prevalence and severity of dental caries in schoolchildren from the city of Florianópolis, SC, Brazil, in 2009 and to compare these results with data from previous studies carried out in the same institution since 1971. **Methods:** All 145 12- and 13-year-old schoolchildren enrolled in a public school were eligible for the study. Dental caries was assessed with the DMFS/DMFT indexes, the significant caries index (SiC) and their 95% confidence intervals (CI 95%). **Results:** The response rate was 68.2%. The DMFT ranged from 9.2 in 1971 to 0.78 in 2009, although different diagnostic criteria were adopted to estimate these counts. The prevalence of dental caries declined from 98.0% (CI 95% 96.0-100.0) in 1971 to 43.5% (CI 95% 33.7-53.4) in 2009. There was a reduction of 47% in the mean SiC index between 2002 and 2009, i.e. from 3.4 (CI 95% 3.0-3.8) to 1.8 (CI 95% 1.5-2.1). The mean DMFS index was 2.0, and it was composed of an average of 1.8 and 0.2 decayed and filled surfaces, respectively. **Conclusions:** Even though different diagnostic criteria were adopted during the study period, an effective decline in the prevalence and severity of dental caries in adolescents was observed throughout 38 years of monitoring.

Keywords: dental caries, DMF index, oral health, trends.

Introduction

The World Health Organization (WHO) holds epidemiological data showing a decline in the levels of caries experience in 12-year-old children in several countries in recent decades, including some from South America, such as Argentina, Chile, Uruguay, Paraguay, and Brazil. According to the WHO¹, in 2004, the global mean DMFT (an index measuring counts of decayed, missing, and filled permanent teeth) was 1.61. In Brazil, the mean DMFT was 2.8 in 2002-2003, although almost 70% of the Brazilian children aged 12 years and nearly 90% of adolescents aged between 15-19 years presented at least one permanent tooth with dental caries experience, according to the Brazilian Ministry of Health² and the WHO¹.

A bibliographic search carried out in PubMed and Scielo using “dental caries” and “trends” as descriptors, limited to the period between 2004 and 2009, showed that several studies in Brazil documented declining trends in the mean DMFT in 12-year-old adolescents. In the city of Bauru, SP, Bastos et al.³ observed that the dental caries decline was of 84.5% between 1976 and 2001. Gushi et al.⁴, using

Received for publication: May 19, 2010

Accepted: August 24, 2010

Correspondence to:

Marco Aurélio Peres

Departamento de Saúde Pública, Centro de Ciências da Saúde

Universidade Federal de Santa Catarina

Campus Universitário – Trindade - 88010-970

Florianópolis - SC

Phone: +55-48-37219046

Fax: +55-48-37219542

E-mail: mperes@ccs.ufsc.br

data from oral health surveys carried out with adolescents between 12 and 18 years of age in the state of São Paulo, observed a reduction of 32.3% in the mean DMFT between 1998 and 2002. In the city of Palhoça, SC, Traebert et al.⁵ observed a reduction of 41.9% in the mean DMFT index for adolescents in the same age group between 1997 and 2003. In the city of Goiânia, GO, the decline in dental caries was even greater, reaching 70.6% between 1988 and 2003, as Reis et al.⁶ reported.

Studies carried out in the same school in 1971-1997⁷, 2002⁸, and 2005⁹ in Florianópolis, SC, also demonstrated an expressive reduction in the severity (91.3%) and in the prevalence (58.2%) of dental caries. Worldwide, very few studies have monitored dental caries prevalence and severity in the same population group for a long period. In Australia, Armfield et al.¹⁰ conducted a study with schoolchildren enrolled in public and private schools from 1977 to 2002 and observed a reduction of 79% in the experience of dental caries throughout 25 years. In Somerset, United Kingdom, Anderson¹¹ reported that the mean DMFT declined by 80% in 12-year-old adolescents from two schools, between 1963 and 1988.

The monitoring of dental caries over time is a part of oral health surveillance system; it is important to subsidize health planners to implement and evaluate preventive programs and dental care. The aim of this study was to evaluate the prevalence and severity of dental caries in 12- and 13-year-old schoolchildren from Florianópolis, SC, Brazil, in 2009 and to compare these data to the results obtained in previous studies conducted in the same school in 1971, 1997, 2002, and 2005.

Material and methods

The sample comprised all 145 students 12 and 13 years of age, enrolled in a public school (Escola de Educação Básica Padre Anchieta) in the city of Florianópolis, state of Santa Catarina, Southern Brazil, in 2009. The same methodological procedures used in previous studies⁷⁻⁹ were adopted in the present investigation, including the collection of clinical data under natural light and using plane dental mirrors and periodontal probes (ball-end probes), following the diagnostic criteria recommended by the WHO¹².

The fieldwork team was composed of two undergraduate dental students; the first performed all dental examinations, whereas the second transferred clinical records to appropriate forms. Both were supervised and trained by an experienced dental epidemiologist (MAP) to ensure standardization of dental caries diagnosis, as well as to avoid discrepancies in interpretation of dental examinations. This calibration training consisted of repeated dental examinations on 15 subjects, followed by discussion on the dental caries diagnostics. This process was ended when the discrepancies were eliminated, and consistency in dental caries diagnosis was achieved among dental examiners. For this phase of the fieldwork, intra- and interexaminer reliabilities were not estimated.

Because a remarkable decline in dental caries prevalence

over the past decades has been observed, the DMFS index (number of decayed, missing, or filled permanent dental surfaces) was used as a diagnostic criterion, in addition to the DMFT index. A tooth surface was considered as "decayed" when a lesion in a pit or fissure or a smooth tooth surface presented an unmistakable cavity, undermined enamel, or a detectable softened floor or walls. A filled but still decayed surface was also included in this category. This index, recommended by the WHO¹², also considered further areas with past experience of dental caries, recording those that received restorative dental treatment and those that have been extracted because of caries.

According to the methodology proposed by Pinto¹³, the 28 permanent teeth were divided into 128 surfaces, as follows: molars and premolars with five surfaces (occlusal, mesial, distal, buccal, and lingual) and incisors and canines with four surfaces (mesial, distal, vestibular, and lingual). To record clinical data, the following sequence was respected: upper right (2nd molar to the central incisor), upper left (central incisor to 2nd molar), lower left (2nd molar to central incisor), and lower right (central incisor to 2nd molar). To analyze tooth surfaces, the examination started in the occlusal surface (O), followed by mesial (M), buccal (B), distal (D), and lingual (L) surfaces.

Data were typed into a Microsoft® Excel spreadsheet. Statistical analysis was performed with Stata, v.9. The prevalence of dental caries, the mean DMFS index, the mean DMFT index, and their components were also estimated with 95% confidence intervals. The chi-square test was used to compare caries prevalence according to sex, and the Mann-Whitney U test was used to compare the distributions of DMFS according to sex and age. The significant caries index (SiC) was calculated based on the mean DMFT index of the highest DMFT tertile in the studied population¹⁴.

A letter was sent to parents explaining the aims and importance of the study and asking permission for their children's participation. When the survey was completed, another letter was sent to acknowledge the participation of adolescents, to report any dental treatment needs, and to provide advices on oral health. The project was approved by the Ethics Committee for Human Research of the Federal University of Santa Catarina (protocol number 126249/2008-3).

Results

From a total of 148 eligible schoolchildren, 101 took part in the study (response rate of 68.2%). Losses were due to refusals (18.2%) and transferences to other schools, during data collection period. The distribution of the studied adolescents by age and sex in the present and previous studies is shown in Table 1. In 1971, 1997, 2002, and 2005, most of the schoolchildren were aged 12 years; this proportion ranged from 62.9% in 1971 to 45.5% in 2009. However, in 2009, 13-year-old children were more frequent (54.5%), when taking the 1971 study as a reference.

The differences in caries prevalence by sex and age were not statistically significant ($p = 0.4$ and $p = 0.1$, respectively). Consequently, the results are shown for the sample as a whole.

Table 1. Distribution by age and sex of 12- and 13-year-old schoolchildren enrolled in Escola de Educação Básica Padre Anchieta in 1971, 1997, 2002, 2005 and 2009. Florianópolis, Santa Catarina, Brazil, 2009.

Variables	1971		1997		2002		2005		2009	
	N	%	N	%	N	%	N	%	N	%
Age										
12 years	127	62.9	101	57.7	88	52.1	118	69.0	46	45.5
13 years	75	37.1	74	42.3	81	47.9	53	31.0	55	54.5
Gender										
Male	*	*	90	51.4	77	45.6	81	47.4	48	47.5
Female	*	*	85	48.6	92	54.4	90	52.6	53	52.5
Total	202	100.0	175	100.0	169	100.0	171	100.0	101	100.0

* Unavailable information

Table 2 displays the mean DMFT and its components, as well as 95% confidence intervals. In the present study, the mean D component was 0.65 (CI 95% 0.45-0.83), followed by the F component with 0.12 (CI 95% 0.03-0.22) and by the M component with 0.01 (CI 95% 0.00- 0.03). The mean DMFT index was 0.78 (CI 95% 0.56-1.00). In the later period of the present study (2009), the D component contributed most to the mean DMFT value, corresponding to 83.3% of the index. When compared to the percent composition of the DMFT index in the first study year (1971), there was a greater reduction of the M component (97.43%), followed by the D (91.71%) and F (87.09%) components in the present study.

Caries prevalence, i.e. the number of participants with DMFT¹ divided by all examined schoolchildren, declined from 98.0% (CI 95% 96.0-100.0) in 1971 to 93.7% (CI 95% 90.0-98.0) in 1997, when the Klein and Palmer (1938) diagnostic criteria were adopted. Between 1997 and 2002, and taking into account the WHO 1987 and the WHO 1997 diagnostic criteria, respectively, the caries prevalence decreased from 80.0% (CI 95% 74.0-86.0) to 57.4% (CI 95% 50.0-65.0). In the penultimate period of observation (2002-2005), caries prevalence declined from 57.4% (CI 95% 50.0-65.0) to 40.9% (CI 95% 33.5-48.4), according to the criteria set forth by the WHO¹² in 1997. In the later period of the

present study (2009), the dental caries prevalence reached 43.5% (CI 95% 33.7- 53.4).

The average of DMFT index was 9.2 (CI 95% not available) in 1971, declining to 6.2 (CI 95% 5.2-7.4) in 1997, according to the Klein and Palmer (1938) diagnostic criteria. The mean reached 3.0 (CI 95% 2.7-3.3) and 1.4 (CI 95% 1.1-1.6) in 2002, when the WHO 1987 and the WHO 1997 diagnostic criteria were adopted, respectively. In 2005, the mean DMFT index was 0.8 (CI 95% 0.6-1.0), a value also observed in the present study (Figure 1).

In 2002, the SiC index was 3.4 (CI 95% 3.0-3.8), decreasing to 2.2 (CI 95% 1.8- 2.5) in 2005 and to 1.8 (CI 95% 1.5-2.1) in the present study. This corresponds to a reduction of 47% (Figure 2). Table 3 shows that the mean decayed and filled surfaces in 2009 was 1.8 (CI 95% 1.1-2.5) and 0.2 (CI 95% 0.0-0.4), respectively. Seventy-five percent (75th percentile) of the adolescents showed a mean of decayed and filled surfaces (D and F) below 2. In 2009, the mean DMFS index was found to be 2.0, with an average of decayed surfaces of 2.3 in males and 1.7 in females, a nonsignificant difference. At the age of 13 years, an average of 2.7 decayed and filled surfaces was observed; at the age of 12 years, this average was 1.2. However, this difference was not statistically significant.

Table 2. Mean DMFT Index and its components decayed (D), missing (M) and filled (F) teeth in 12- and 13-year-old schoolchildren enrolled in Padre Anchieta Basic School in 1971, 1997, 2002, 2005 and 2009. Florianópolis, SC, Brazil, 2009.

Year	D			M			F			DMF-T		
	%	mean	95% CI	%	mean	95% CI	%	mean	95% CI	%	mean	95% CI
1971 ¹	85.7	7.85	*	4.2	0.39	*	10.1	0.93	*	100.0	9.17	*
1997 ¹	90.7	5.67	4.57-6.77	0.5	0.03	0.02-0.04	8.8	0.55	0.22-0.88	100.0	6.25	5.15-7.35
1997 ²		2.42									3.00	2.67-3.33
2002 ³	46.8	0.65	0.63-0.67	5.7	0.08	0.03-0.10	47.5	0.66	0.46-0.86	100.0	1.39	1.14-1.64
2005 ³	56.8	0.50	0.30-0.60	4.9	0.04	0.01-0.06	38.3	0.3	0.20-0.40	100.0	0.84	0.60-1.00
2009 ³	83.3	0.65	0.45-0.83	1.3	0.01	0.00- 0.03	15.4	0.12	0.03-0.22	100.0	0.78	0.56-1.00

*Unavailable information

¹Klein and Palmer (1938)²¹ diagnostic criteria²WHO diagnostic criteria (1987)²²³WHO diagnostic criteria (1997)¹²

Table 3. Descriptive statistics of decayed and filled surfaces of 12- and 13-year-old schoolchildren enrolled in Escola de Educação Básica Padre Anchieta, Florianópolis Santa Catarina, Brazil, 2009.

	Decayed Surfaces (Ds)	Filled Surfaces (Fs)	Decayed and Filled Surfaces (DFs)
Mean	1.8	0.2	2.0
CI 95%	1.1-2.5	0.0-0.4	1.2-2.8
Median	0.0	0.0	0.0
Minimum	0	0	0
Maximum	20	8	20
Percentile 75	1.5	0.0	2.0




Fig.1 - Mean DMFT index and its decayed, missing, and filled components in the years 1971, 1997, 2002, 2005, and 2009 of 12- and 13-year-old schoolchildren enrolled in Escola de Educação Básica Padre Anchieta in 1971, 1997, 2002, 2005, and 2009. Florianópolis, Santa Catarina, Brazil, 2009.

* Unavailable information; ¹Klein and Palmer (1938)²¹ diagnostic criteria
²WHO diagnostic criteria (1987)²²; ³WHO diagnostic criteria (1997)¹²




Fig. 2 - Trend of mean Significant Caries Index (SiC) of 12- and 13-year-old schoolchildren enrolled in Escola de Educação Básica Padre Anchieta from 2002 to 2009. Florianópolis, Santa Catarina, Brazil, 2009.

Discussion

Since 1971, when the first of this series of studies was conducted, until 2009, an effective decline in dental caries prevalence and severity was observed in the studied population. Even though different diagnostic criteria were used along the studied period, a reduction of 91.4% in the caries severity was found in the 2009 study, when compared with the 1971 study. Compared to the 2002 study, which adopted the same diagnostic criteria, a reduction of 43.4% in the caries severity was observed in 2009.

Despite the fact that the mean DMFT did not change between 2005 and 2009, its components presented contradictory proportions when compared with those observed in previous studies. The D component showed a reduction of 93.6% between 1971 and 2005, represented 83.3% of the

DMFT in 2009, and showed a relative increase of 30% over the last studied period (2005-2009). These data may hypothetically indicate a decrease in dental care utilization in the region, a paradox considering that, for this period, the number of dentists in the city increased, achieving a ratio of 1 dentist for every 250 inhabitants in the city of Florianópolis in June 2009. A less invasive approach by dentists in the city may be another explanation for such a decline in the proportion of D component.

The 2009 F component showed a reduction of 60% as compared with the previous study (2005) and comprised 15.4% of the DMFT index in 2009. The M component showed an even greater reduction (75%) since the last year of study, although it represents only 1.3% of the mean DMFT index in the current study. The mean DMFT was 0.78, reaching one of the WHO goals for the year 2010, that is, a mean DMFT below 1.0 for the age of 12 years, according to Gomes et al.¹⁵.

It is noteworthy that there are few studies conducted in the same place for such a long period and under the supervision of the same research team. Marcenes et al.¹⁶ and Souza et al.⁹, suggested that the reduction observed in dental caries experience may be partially attributed to changes in diagnostic criteria for dental caries over time.

In Brazil, similar decline in dental caries was demonstrated by other researchers. As stated by Martins et al.¹⁷ the city of in Bilac, SP, a similar reduction in the mean DMFT index was achieved, despite the lack of water fluoridation supply during the study period. According to Bastos et al.³ (2005), fluoridated water supplies contributed to the remarkable reduction in the mean DMFT in Bauru, SP. In fact, the results from the "SB Brazil: Oral Health in the Brazilian Population" held in 2003, produced by Ministry of Health², revealed that in the South region - an area with high water fluoridated coverage - presented 37.7% of caries-free adolescents aged 12 years, in contrast to only 12.9% of caries-free adolescents in non-fluoridated regions.

Worldwide, similar reductions in caries experience have been documented. Armfield and Spencer¹⁰ (2008) analyzed data from dental caries in adolescents enrolled in public and private schools between 1977 and 2002 in Australia and found a reduction of 79% for the age of 12 years. In this same study, an annual reduction of 2.6% the SiC index between 1977 and 2002 for the same age was observed. These values were lower than the ones found in the present study (annual reduction of 6.7%). In Lithuania (1993-2001), Aleksejūnienė et al.¹⁸ (2004) reported that the decline in dental caries rates may be explained by an improvement in oral hygiene because

products for this purpose began to be sold in 1991 and had an increase of 95% in their use in 2001 in that country.

The US Centers for Diseases Control and Prevention¹⁹ declared that the fluoridation of water supplies is a the most effective mass intervention to prevent dental caries, and it was considered one of the most important public health achievements in the last century. In addition to water fluoridation, the major decline in dental caries prevalence and severity has been attributed to the access to other fluoride sources, especially to the widespread use of fluoridated toothpaste. Moreover, an improvement in socioeconomic indicators and the boost of initiatives in dental public health also have been listed, by Bastos et al.³, Ministry of Health²⁰, Martins et al.¹⁷, as important to dental caries reduction over time. As stated by Bastos et al.³ and Martins et al.¹⁷, the use of fluoridated toothpaste in Brazil was intensified from 1989, when the Ministry of Health regulated the addition of fluoride in these products and mouthrinses. Possibly, according to Freysleben et al.⁷, Bastos et al.⁸ and Souza et al.⁹, these factors contributed to a higher rate of annual reduction in the dental caries severity over the periods that followed the studies through the 1990s in Florianópolis.

However, the design of this study does not allow an assessment of the possible reasons for the decline in dental caries prevalence. In the first period (1971 to 1997), in which the lowest annual reduction rate was observed compared with other years, this could be explained, as noted by Freysleben et al.⁷, by the shorter period of public exposure to fluoridated water, implemented in Florianópolis in 1982, and the availability of use of fluoride toothpaste from the late 1980's. In 1971, the school was located in a suburban area of the city, which became, in 1997, a middle class region with a residential profile and close to the city's downtown with better living conditions. In the second studied period (1997 to 2002), the caries decline can be attributed to the increased availability of fluoridated toothpaste and to the implementation of fluoridated water supply. Additionally, Florianópolis in 2003 occupied the third position among the capitals with the lowest level of social exclusion in Brazil. In 2008, Bastos et al.⁸ reported that Florianópolis appeared as the city with the highest human development index (HDI of 0.86) among 33 other Brazilian metropolitan areas and cities and presented the highest average income among Brazilian cities. Despite the observed declining trend in dental caries over the studied period, different authors^{10-11,18} found that the disease rate is likely to stabilize because of the low value already reached.

Some limitations of this study must be mentioned. The 2009 study had a relatively low response rate, lower than those observed in the previous phases. We can hypothesize that the mean DMFT could be higher, assuming that individuals who were not examined presented higher dental caries attack than those who effectively took part in the study.

In conclusion, despite the fact that different dental caries diagnostic criteria were used, there was an actual decrease in the mean DMFT over the 38-year study period. It should be noted that these results cannot be readily generalized to the entire population of 12 and 13 years of age from Florianópolis, SC.

Acknowledgments

Helena Mendes Constante was supported by a scholarship granted from the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) of Brazil from November 2008 to July 2009 (number 126249/2008 3).

References

- World Health Organization. Oral health country profiles. Available from: <http://www.whocollab.od.mah.se/index.html>. Accessed 2009 Aug.
- Brasil. Ministério da Saúde. Secretaria de Atenção à Saúde. Departamento de Atenção Básica. Projeto SB Brasil 2003: condições de saúde bucal da população brasileira 2002-2003: resultados principais. Brasília: Ministério da Saúde; 2004.
- Bastos RS, Olympio KPK, Bijella VT, Buzalaf MAR, Bastos JRM. Trends in dental caries prevalence in 12-year-old schoolchildren between 1976 and 2001 in Bauru, Brazil. *Public Health*. 2005; 119: 269-75.
- Gushi LL, Rihs LB, Soares MC, Forni TIB, Vieira V, Wada RS, et al. Cárie dentária e necessidades de tratamento em adolescentes do estado de São Paulo, 1998 e 2002. *Rev Saúde Pública*. 2008; 42: 480-6.
- Traebert J, Lacerda JT, Fischer TK, Junbo Y. Dental Caries and Orofacial Pain Trends in 12-year-old school children between 1997 and 2003. *Oral Health Prev Dent*. 2005; 3: 243-8.
- Reis SCGB, Freire MCM, Higino MASP, Batista SMO. Declínio de cárie em escolares de 12 anos da rede pública de Goiânia, Goiás, Brasil, no período de 1988 a 2003. *Rev Bras Epidemiol*. 2009; 12, 1: 92-8.
- Freysleben GR, Peres MAA, Mercenes W. Prevalência de cárie e CPO-D médio em escolares de doze a treze anos de idade nos anos de 1971 e 1997, região Sul, Brasil. *Rev Saúde Pública*. 2000; 34: 304-8.
- Bastos JLD, Nomura LH, Peres MA. Trends in dental caries rates in schoolchildren 12 and 13 years old in Florianópolis, Santa Catarina, Brazil, 1971-2002. *Cad Saúde Pública*. 2004; 20: 117-22.
- Souza ML, Bastos JLD, Peres MA. Trends in dental caries rates in 12- and 13-year old schoolchildren from Florianópolis (Brazil) between 1971 and 2005. *Oral Health Prev Dent*. 2006; 4: 187-92.
- Armfield JM, Spencer AJ. Quarter of a century of change: caries experience in Australian children, 1977-2002. *Aust Dent J*. 2008; 53: 151-9.
- Anderson RJ. The changes in dental caries experience of 12-year-old schoolchildren in two Somerset schools. A review after an interval 25 years. *Br Dent J*. 1989; 167: 321-4.
- World Health Organization. Oral health surveys: basic methods. 4th ed. Geneva: World Health Organization; 1997.
- Pinto VG. Saúde bucal e coletiva. 4. ed. São Paulo: Santos; 2000.
- Bratthall D. Introducing the Significant Caries Index together with a proposal for a new global oral health goal for 12-year olds. *Int Dent J*. 2000; 50: 378-84.
- Gomes PR, Costa SC, Cypriano S, Sousa MLRP, São Paulo, Brasil: situação da cárie dentária com relação às metas OMS 2000 e 2010. *Cad Saúde Pública* 2004; 20: 866-70.
- Marcenes W, Freysleben GR, Peres MA. Contribution of changing diagnostic criteria toward reduction of caries between 1971 and 1997 in children attending the same school in Florianópolis, Brazil. *Community Dent Oral Epidemiol* 2001; 29: 449-55.
- Martins RJ, Garbin CAS, Garbin AJI, Moimaz SAS, Saliba O. Declínio de cárie em um município da região noroeste do Estado de São Paulo, Brasil, no período de 1998 a 2004. *Cad Saúde Pública*. 2006; 22: 1035-41.
- Aleksejūnienė J, Holst D, Balciūnienė I. Factors influencing the caries decline in Lithuanian adolescents—trends in the period 1993-2001. *Eur J Oral Sci*. 2004; 112: 3-7
- Centers for Diseases Control and Prevention (1999). Achievements in public health, 1900-1999. Fluoridation of drinking water to prevent dental caries. *Morb Mortal Wkly Rep*. 44(RR-13): 1-40.
- Brasil. Ministério da Saúde. Secretaria Nacional de Vigilância Sanitária. Portaria n. 22, de 20/12/1989. Diário Oficial União, Seção I. Available from: http://dtr2004.saude.gov.br/dab/docs/legislacao/portaria22_20_12_89.pdf. Accessed 2009 Aug.
- Klein H, Palmer CE. Dental caries in American Indian children. Washington DC: Government Printing; 1938.
- World Health Organization. Oral health surveys: basic methods. 3rd ed. Geneva: World Health Organization; 1987.